

## B. ANALYSIS OF PERFORMANCE

### CHEMISTRY PAPER 1 (THEORY)

#### PART I (20 Marks)

Answer *all* questions.

#### Question 1

- (a) Fill in the blanks by choosing the appropriate word/words from those given in the brackets: [5]  
(zero, first, second, increased, decreased, anode, cathode, active, inactive, potassium cyanide, internal, external, dependent, independent, red, benzoic acid, benzoin, common ion effect, salt hydrolysis, alkali, potassium hydroxide.)
- (i) In a galvanic cell, electrons flow from \_\_\_\_\_ to \_\_\_\_\_ through the connecting wires.
- (ii) Racemic mixtures are optically \_\_\_\_\_ because of \_\_\_\_\_ compensation.
- (iii) Half life period of a \_\_\_\_\_ order reaction is \_\_\_\_\_ of the concentration of the reactant.
- (iv) Benzaldehyde when treated with an alcoholic solution of \_\_\_\_\_ forms \_\_\_\_\_.
- (v) Solubility of calcium oxalate is \_\_\_\_\_ in the presence of ammonium oxalate because of \_\_\_\_\_.
- (b) Complete the following statements by selecting the **correct alternative** from the choices given: [5]
- (i) The compound which is optically active is:
- (1) 1-butanol
  - (2) 2-butanol
  - (3) 1-propanol
  - (4) 2-methyl-1-propanol
- (ii) The salt which will not hydrolyse in aqueous solution is:
- (1) Copper sulphate
  - (2) Sodium sulphate
  - (3) Potassium cyanide
  - (4) Sodium carbonate
- (iii) Copper has the face centred cubic structure. The coordination number of each ion is:
- (1) 4
  - (2) 12

(3) 14

(4) 8

(iv) For the reaction  $2\text{SO}_2 + \text{O}_2 \rightleftharpoons 2\text{SO}_3$ , the unit of equilibrium constant is:

(1)  $\text{L mol}^{-1}$

(2)  $\text{J mol}^{-1}$

(3)  $\text{mol L}^{-1}$

(4)  $[\text{L mol}^{-1}]^2$

(v) The deficiency of vitamin D causes:

(1) Rickets

(2) Gout

(3) Scurvy

(4) Night blindness.

(c) Answer the following questions:

[5]

(i) Two metallic elements A and B have the following standard oxidation potentials:

$A = 0.40\text{v}$   $B = -0.80\text{v}$ . What would you expect if element A was added to an aqueous salt solution of element B? Give a reason for your answer.

(ii) Two moles of  $\text{NH}_3$  are introduced into one litre flask in which it dissociates at high temperature as follows:  $2\text{NH}_3(\text{g}) \rightleftharpoons \text{N}_2(\text{g}) + 3\text{H}_2(\text{g})$ . Determine Kc, if at equilibrium 1 mole of  $\text{NH}_3$  remains.

(iii) Give balanced equation for the preparation of salicylaldehyde from phenol.

(iv) If the half life period for a first order reaction is 69.3 seconds, what is the value of its rate constant?

(v) Define cryoscopic constant.

(d) Match the following:

[5]

(i) Colligative property

(a) Polysaccharide

(ii) Nicol prism

(b) Osmotic pressure

(iii) Activation energy

(c) Aldol condensation

(iv) Starch

(d) Polarimeter

(v) Acetaldehyde

(e) Arrhenius equation

### Comments of Examiners

- (a) (i) Some candidates wrote 'cathode to anode'.  
(ii) Many candidates wrote 'active' and 'internal' in place of the correct words.  
(iii) Some candidates wrote 'dependent' in place of 'independent'.  
(iv) Many candidates wrote 'potassium hydroxide' and 'Benzoic acid' instead of the correct words.  
(v) A few candidates wrote 'increased' in place of 'decreased'.
- (b) (i) '1-butanol' or 2-methyl propanol was chosen by some candidates.  
(ii) 'Potassium cyanide' or 'sodium carbonate' was chosen in some cases.  
(iii) Some candidates wrote '4' in place of the correct option.  
(iv) Many candidates chose the options (3) and (4).  
(v) A few candidates chose 'scurvy' or 'night blindness'.
- (c) (i) Candidates were confused with oxidation and reduction potential. Some wrote 'B displaced A' and various other wrong answers and incorrect reasons.  
(ii) Most of the candidates calculated  $K_e$  for 2 moles of  $\text{NH}_3$ . Unit was wrong in many scripts.  
(iii) Wrong reagents were used by many candidates. In some cases, the equation was unbalanced.  
(iv) Several candidates used a wrong formula.  
(v) Most of the candidates gave the mathematical expression and explained that.
- (d) Most of the candidates were able to attempt this part correctly.

### Suggestions for teachers

- The concept of production of current and flow of electrons in a galvanic cell should be explained clearly.
- The idea of 'internal' and 'external' compensation should be taught with correct examples.
- Reactions should be practiced in class with correct names of the reactants and products.
- 'Common ion effect' should be taught in class with equations showing the shift of equilibrium.
- The concept of chiral carbon atom should be made clear. Isomerism should be practised with examples.
- The units of  $K_e$  should be derived in class taking examples.
- The use of electrochemical series should be taught in details giving its importance in determining the properties and nature of the elements.
- Reactions should be practiced with correct reactions and products in the balanced form.

### **MARKING SCHEME**

#### **Question 1.**

- (a) (i) anode, cathode  
(ii) inactive, external  
(iii) first \ 1<sup>st</sup>, independent or zero, dependent or second dependent  
(iv) potassium cyanide, benzion/ formula  
(v) decreased, common ion effect
- (b) (i) 2 butanol / formula  
(ii) (1) Sodium sulphate / formula

(iii) (2) 12 /twelve

(iv) (1) L mol<sup>-1</sup>

(v) (1) Rickets

Answer the following questions:

(c) (i) A displaces B from its salt solution. Standard reduction potential of A is less than standard reduction potential of B or any other explanation.

(ii)  $2\text{NH}_3 \rightleftharpoons \text{N}_2 + 3\text{H}_2$

Initial 2            0    0


At eqn. 1             $\frac{1}{2}$      $\frac{3}{2}$

$[\text{NH}_3] = \frac{1}{1}$  moles per litre     $[\text{N}_2] = \frac{1}{2}$  moles per litre

$[\text{H}_2] = \frac{3}{2}$  moles per litre

$$K_c = \frac{[\text{N}_2][\text{H}_2]^3}{[\text{NH}_3]^2} = \frac{\left(\frac{1}{2}\right)\left(\frac{3}{2}\right)^3}{(1)^2} \text{ mole}^4 / \text{litre}^4$$
$$= \frac{27}{16} \text{ mole}^2 / \text{litre}^2 = 1.687 (\text{mol} / \text{lit})^2$$

(iii)



(iv)  $K = \frac{0.693}{693} = 0.01 \text{sec}^{-1}$

(v) Depression of freezing point when the molality of the solution is unity / any correct definition.

(d) Match the following:

(i) (b) Osmotic pressure

(ii) (d) Polarimeter

(iii) (e) Arrhenius equation

(iv) (a) Polysaccharide

(v) (c) Aldol condensation

## PART II (50 Marks)

Answer six questions choosing two from Section A, two from Section B and two from Section C.

### SECTION A

Answer any two questions.

#### Question 2

- (a) (i) Ethylene glycol is used as an antifreeze agent. Calculate the amount of ethylene glycol to be added to 4 kg of water to prevent it from freezing at  $-6^{\circ}\text{C}$ . ( $K_f$  for  $\text{H}_2\text{O} = 1.85 \text{ K mole}^{-1} \text{ kg}$ ) [3]
- (ii) The freezing point of a solution containing 0.3gms of acetic acid in 30gms of benzene is lowered by  $0.45\text{K}$ . Calculate the Van't Hoff factor. (at. wt. of C = 12, H = 1, O = 16,  $K_f$  for benzene =  $5.12\text{K kg mole}^{-1}$ ). [2]
- (b) Name the law or principle confirmed by the following observations: [2]
- (i) When water is added to  $0.01\text{M}$  aqueous solution of acetic acid the number of hydrogen ions increase.
- (ii) When  $96500$  coulombs of electricity is passed through acidulated water,  $5.6$  litres of oxygen at s.t.p. is liberated at the anode.
- (c) Arrange Ag, Cr and Hg metals in the increasing order of reducing power. Given: [1]
- $$E^{\circ}_{\text{Ag}^+/\text{Ag}} = +0.80\text{V}$$
- $$E^{\circ}_{\text{Cr}^{3+}/\text{Cr}} = -0.74\text{V}$$
- $$E^{\circ}_{\text{Hg}^{2+}/\text{Hg}} = +0.79\text{V}$$
- (d) In a first order reaction, 10% of the reactant is consumed in 25 minutes. Calculate: [2]
- (i) The half life of the reaction.
- (ii) The time required for completing 17% of the reaction.

#### Comments of Examiners

- (a) (i) Incorrect formula and substitution was done by many candidates. Some candidates were unable to calculate the molecular weight of glycol.
- (ii) Many candidates made wrong calculations due to wrong substitution.
- (b) (i) Many candidates gave the answer as, 'dilution law' instead of Ostwald dilution Law.
- (ii) Some candidates wrote only 'Faraday's law' but did not mention the '2<sup>nd</sup> law'.

#### Suggestions for teachers

- More practice should be given in numericals along with correct substitution.
- Important laws and their applications should be taught in class with correct examples.
- Electrochemical series should be taught with reference to the properties of the elements.

- (c) Several candidates were confused between 'reduction' and 'reduction potential'. Many candidates arranged the metals in the reverse direction.
- (d) (i) Some candidates made calculation errors in this part of the question.  
(ii) Incorrect formula was taken by some candidates.

### MARKING SCHEME

#### Question 2.

(a) (i) 
$$\Delta T_f = K_f \times \frac{w \times 1000}{m \times W}$$

$$w = \frac{\Delta T_f \times m \times W}{K_f \times 1000} \quad [\text{Mol. wt. of } \begin{array}{c} \text{CH}_2\text{OH} \\ | \\ \text{CH}_2\text{OH} \end{array} = 62 ]$$

$$= \frac{6 \times 62 \times 4 \times 1000}{1.85 \times 1000}$$

$$= 804.32 \text{ g}$$

(ii) 
$$\Delta T_f = iK_f \times \frac{w \times 1000}{m \times W} \quad \text{or} \quad T_f = i k_f \times m$$

$$i = \frac{\Delta T_f \times m \times W}{K_f \times w \times 1000}$$

$$= \frac{0.45 \times 60 \times 30}{5.12 \times 0.3 \times 1000}$$

$$= 0.527$$

(b) (i) Ostwald's dilution law

(ii) Faraday's second law

(c) Ag, Hg, Cr

(d) 
$$K = \frac{2.303}{t} \log \left( \frac{a}{a-x} \right)$$

$$K = \frac{2.303}{25} \log \left( \frac{100}{90} \right)$$

$$= 4.215 \times 10^{-3} \text{ min}^{-1}$$

$$\text{Half life } (t_{1/2}) = \frac{0.693}{K} = \frac{0.693}{4.215 \times 10^{-3}} = 164.41 \text{ minutes}$$

$$t = \frac{2.303}{k} \log \left( \frac{a}{a-x} \right)$$

$$= \frac{2.303}{4.215 \times 10^{-3}} \log \left( \frac{100}{83} \right) = 44.21 \text{ minutes}$$

### Question 3

- (a) Explain giving reasons why (Give equations in support of your answer):
- (i) A solution of  $\text{NH}_4\text{Cl}$  and  $\text{NH}_4\text{OH}$  acts as a buffer. [2]
- (ii) Cu is precipitated as  $\text{CuS}$  while Zn is not precipitated when  $\text{H}_2\text{S}$  is passed through an acidic solution of  $\text{Cu}(\text{NO}_3)_2$  and  $\text{Zn}(\text{NO}_3)_2$  respectively. [2]
- (b) (i) What is Schottky defect in a solid? [1]
- (ii) A bcc element (atomic mass 65) has a cell edge of 420 pm. Calculate its density in  $\text{gms/cm}^3$ . [3]
- (c) The rate of the reaction  $\text{H}_2 + \text{I}_2 \rightleftharpoons 2\text{HI}$  is given by: [2]

$$\text{rate} = 1.7 \times 10^{-19} [\text{H}_2][\text{I}_2] \text{ at } 25^\circ\text{C}.$$

The rate of decomposition of gaseous HI to  $\text{H}_2$  and  $\text{I}_2$  is given by:

$$\text{rate} = 2.4 \times 10^{-21} [\text{HI}]^2 \text{ at } 25^\circ\text{C}.$$

Calculate the equilibrium constant for the formation of HI from  $\text{H}_2$  and  $\text{I}_2$  at  $25^\circ\text{C}$ .

#### Comments of Examiners

- (a) (i) The definition of Buffer solution was given by many candidates instead of explaining 'Buffer action'.
- (ii) Many candidates did not mention the low  $K_{\text{sp}}$  for  $\text{CuS}$  and high  $K_{\text{sp}}$  for  $\text{ZnS}$ . Some candidates did not mention the suppression of concentration of Sulphide ion in acidic medium.
- (b) (i) Many candidates did not mention the missing of both anion and cation in ionic crystals. Some candidates confused it with Frenkel defect.
- (ii) Several candidates did not attempt this numerical, while some others did the problem with incorrect formula. In some cases, the value of 'Z' was taken wrongly.
- (c) Value of  $K_c$  was taken as  $\text{rate}_1 / \text{rate}_2$ , giving wrong answers.

#### Suggestions for teachers

- The mechanism of buffer action should be explained with reference to the addition of acid, alkali and dilution. Explanation should be given taking correct equation.
- The use of solubility product in salt analysis should be taught with correct examples. Stress should be laid on key words.
- Defects in crystals should be explained diagrammatically.
- Adequate practice should be given in numericals.

#### MARKING SCHEME

##### Question 3.

- (a) (i)  $\text{NH}_4\text{Cl} \rightleftharpoons \text{NH}_4^+ (\text{High}) + \text{Cl}^- (\text{High})$   
 $\text{NH}_4\text{OH} \rightleftharpoons \text{NH}_4^+ (\text{low}) + \text{OH}^- (\text{low})$   
Addition of HCl,  $\text{HCl} \rightleftharpoons \text{H}^+ + \text{Cl}^-$   
 $\text{H}^+ (\text{from HCl}) + \text{OH}^- (\text{from NH}_4\text{OH}) \rightarrow \text{H}_2\text{O}$   
 $\text{OH}^-$  ions are removed from the solution.

In the absence of  $\text{OH}^-$ ,  $\text{NH}_4\text{OH}$ , dissociates more in giving more  $\text{OH}^-$  ions. These  $\text{OH}^-$  ions combines with  $\text{H}^+$  to form water. Process keeps on happening till all  $\text{H}^+$  ions are consumed and pH remains constant.

Addition of NaOH:



$\text{NH}_4\text{OH}$  being a weak base dissociates partially and whatever  $\text{OH}^-$  are released are not sufficient to increase the pH. OR any other explanation.



In the presence of HCl, the dissociation of  $\text{H}_2\text{S}$  is suppressed due to common  $\text{H}^+$  ion and less  $\text{S}^{2-}$  ions are released in solution.

The  $K_{\text{sp}}$  of CuS is low and hence  $[\text{Cu}^{+2}][\text{S}^{2-}] > K_{\text{sp}}$ .

ZnS has a high value of  $K_{\text{sp}}$  and hence, does not get precipitated with less  $\text{S}^{2-}$  ions.


(b) (i) Schottky defect – pair of holes exist in the crystal lattice due to one positive ion and one negative ion being absent from the crystal lattice. OR any other definition

(ii) Density (P) = 
$$\frac{Z \times M}{N_A \times a^3}$$

$$= \frac{2 \times 65}{6.023 \times 10^{23} \times (420 \times 10^{-10})^3}$$

$$= \frac{2 \times 65}{6.023 \times 10^{23} \times 7.4 \times 10^{-23}}$$

$$= 2.91 \text{ gms/cm}^3$$



#### Question 4

- (a) (i) Give Lewis' definition for acids and bases. [1]
- (ii) The solubility of  $\text{Ag}_2\text{CrO}_4$  at  $25^\circ\text{C}$  is  $8.0 \times 10^{-5}$  moles/litre. Calculate its solubility product. [1]
- (b) (i) Define molar conductance of a solution. State its unit. How is it related to the specific conductance of a solution? [2]



- (ii) Calculate the value of  $E_{\text{cell}}$  at 298K for the following cell: [3]



$$E_{\text{Al}^{3+}/\text{Al}}^{\circ} = -1.66 \text{ volt and } E_{\text{Sn}^{2+}/\text{Sn}}^{\circ} = -0.14 \text{ volt}$$

- (c) (i) Calculate the degree of hydrolysis of 0.2(M) sodium acetate solution. [1]  
(Hydrolysis constant of sodium acetate =  $5.6 \times 10^{-10}$  and ionic product of  $\text{H}_2\text{O} = 10^{-14}$  at  $25^{\circ}\text{C}$ )
- (ii) Explain why high pressure is used in the manufacture of ammonia by Haber's process. State the law or principle used. [2]

#### Comments of Examiners

- (a) (i) Several candidates wrote 'electron acceptor' instead of stating 'electron pair acceptor'.  
(ii) Wrong formula was used by some candidates.
- (b) (i) Many candidates explained the formula of molar conductance in place of defining it. The unit was given wrongly in many scripts.  
(ii) Several candidates calculated  $E_{\text{cell}}^{\circ}$  correctly but the calculation of  $E_{\text{cell}}$  was wrong. Some got the same values for both since they rounded up the value instead of calculating it up to the third place of decimal. Many candidates did wrong substitution in the 'Nernst' equation.
- (c) (i) Incorrect formula was used by many candidates.  
(ii) The explanation given by many candidates was not clear. In some cases, Le Chatelier's principle was only mentioned but not stated.

#### Suggestions for teachers:

- Lewis definition should be correctly taught with correct examples.
- Derivation of the  $K_{\text{sp}}$  values of different salts should be done in class.
- 'Nernst equation' should be taught with correct examples and correct substitution.
- Numericals should be practised with correct formula of substitution.
- Le Chatelier's principle should be explained with reference to the equations mentioned in the scope of syllabus.

#### MARKING SCHEME

##### Question 4.

- (a) (i) Lewis acid is an electron pair acceptor.

Lewis base is an electron pair donor.



$$K_{\text{sp}} = [\text{Ag}^+]^2[\text{CrO}_4^{2-}]$$

$$= (2S)^2(S) = 4S^3$$

$$K_{\text{sp}} = 4 \times (8 \times 10^{-5})^3 = 2.048 \times 10^{-12}$$

- (b) (i) Molar conductance is defined as the conducting power of all the ions produced by dissolving one gm mole of an electrolyte in solution/ any other definition

Its unit is  $\text{ohm}^{-1} \text{cm}^2 \text{mole}^{-1}$  or  $\text{ohm}^{-1} \text{m}^2 \text{mole}^{-1}$

$$\Lambda_m = \frac{1000}{c} \times \Lambda_{sp}$$




c = conc. in molarity

(ii)  $E_{\text{cell}}^{\circ} = (E_{\text{redn}}^{\circ})_{\text{cathode}} - (E_{\text{redn}}^{\circ})_{\text{anode}}$

$$= (-0.14) - (1.66) \text{ volt}$$

$$= 1.52 \text{ volt}$$

$$E_{\text{cell}} = E_{\text{cell}}^{\circ} - \frac{0.05912}{n} \log \frac{[\text{products}]}{[\text{Reactants}]}$$



$$E_{\text{cell}} = 1.52 - \frac{0.05912}{6} \log \frac{[\text{Al}^{3+}]^2 [\text{Sn}]^3}{[\text{Al}]^2 [\text{Sn}^{2+}]^3} \text{ volt}$$

$$= 1.52 - \frac{0.05912}{6} \log \frac{(0.01)^2 (1)^3}{(1)^2 (0.015)^3} \text{ volt}$$

$$= (1.52 - 0.0145) \text{ volt} = 1.5055 \text{ volt}$$

(c) (i)  $h = \sqrt{\frac{kw}{ka.c}} = \sqrt{\frac{Kh}{c}}$

$$= \sqrt{\frac{5.6 \times 10^{-10}}{0.2}} = 5.29 \times 10^{-5}$$

- (ii) Formation of ammonia takes place with decrease in volume. Hence, the reaction is favoured with high pressure.

Le Chatelier's Principle – when a system at equilibrium is subjected to a stress, the equilibrium shifts in that direction to nullify the effect of the stress OR any other explanation.

## SECTION B

Answer any two questions.

### Question 5

- (a) Give the IUPAC names of the following coordination compounds: [2]
- $K_2[Zn(OH)_4]$
  - $[Co(NH_3)_5(CO_3)]Cl$
- (b) For the complex ion  $[Fe(CN)_6]^{3-}$  state: [1]
- The geometry of the ion.
  - The magnetic property of the ion.
- (c) What type of structural isomers are  $[Co(NH_3)_5Br]SO_4$  and  $[Co(NH_3)_5SO_4]Br$ ? Give a chemical test to distinguish the isomers. [2]

### Comments of Examiners

- (a) (i) Several candidates wrote 'hydroxy' in place of 'hydroxo'. The valency of Zinc was not given correctly by a number of candidates.
- (ii) Several candidates made mistakes in writing the correct name. 'Carbonate' was written instead of 'Carbonato'. The oxidation state of cobalt was given incorrectly by some candidates.
- (b) (i) Many candidates wrote 'tetrahedral' or 'bipyramidal'.
- (ii) Some candidates gave the answer as 'diamagnetic'.
- (c) Many candidates wrote 'linkage isomerism'. Chemical test was not given but the ionisation of the two compounds was shown.

### Suggestions for teachers

- Many examples of co-ordination complexes should be practised with correct spelling and correct oxidation state.
- Geometry of co-ordination compounds and ions should be explained by drawing the correct structure.
- The relation of the magnetic property with number of unpaired electrons should be explained by writing the electronic configuration in box diagrams.

### MARKING SCHEME

#### Question 5.

- (a) (i) potassiumtetrahydroxo zincate(II)
- (ii) pentaamminecarbonatocobalt(III) chloride
- (b) (i) Octahedral, Paramagnetic
- (ii) Ionisation isomerism.
- $[Co(NH_3)_5Br]SO_4$  gives white ppt with  $BaCl_2$  solution, but  $[Co(NH_3)_5SO_4]Br$  does not give white ppt. with  $BaCl_2$  solution OR any other test.
- (c) Type of structural isomers and correct chemical test.

### Question 6

- (a) For the molecule  $\text{XeF}_2$  : [2]
- Draw the structure of the molecule indicating the lone pairs.
  - State the hybridisation of the central atom.
  - State the geometry of the molecule.
- (b) Give balanced chemical equations for the following reactions: [3]
- Fluorine treated with dilute sodium hydroxide solution.
  - Hydrogen sulphide treated with concentrated sulphuric acid.
  - Potassium iodide treated with acidified potassium permanganate solution.

#### Comments of Examiners

- (a) (i) Proper orientation of the orbitals around the Xenon atom was not shown by many candidates. Lone pair of electrons was missing in some cases.
- (ii) The hybridisation was given incorrectly by many candidates.
- (iii) Some candidates gave the answer as 'T' shaped. The word 'linear' was missing in many answers.
- (b) In this part, many candidates wrote unbalanced and incorrect equations.


#### Suggestions for teachers

- Structures of the compounds should be explained by drawing and showing the lone pair of electrons.
- The correct hybridisation should be taught by writing the electronic configuration in box diagrams.
- The geometry of the molecules should be explained on the basis of hybridisation.
- Students should be given practice in writing correct balanced equations.

#### MARKING SCHEME

##### Question 6.

(a)



$sp^3d$ , linear.

- (b) (i)  $2\text{F}_2 + 2\text{NaOH} \rightarrow \text{F}_2\text{O} + 2\text{NaF} + \text{H}_2\text{O}$
- (ii)  $\text{H}_2\text{S} + \text{H}_2\text{SO}_4 \rightarrow 2\text{H}_2\text{O} + \text{SO}_2 + \text{S}$
- (iii)  $2\text{KMnO}_4 + 8\text{H}_2\text{SO}_4 + 10\text{KI} \rightarrow 6\text{K}_2\text{SO}_4 + 2\text{MnSO}_4 + 8\text{H}_2\text{O} + 5\text{I}_2$

### Question 7

- (a) In the extraction of zinc from zinc blende: [2]
- Give an equation to show how zinc oxide is converted to zinc.
  - How is impure zinc finally electro-refined?
- (b) Explain why: [3]
- Transition elements form coloured compounds.
  - Interhalogen compounds are more reactive than their constituent elements.
  - $\text{Cu}^+$  is diamagnetic but  $\text{Cu}^{2+}$  is paramagnetic. ( $Z = 29$ )

### Comments of Examiners

- (a) (i) Some candidates wrote 'CO<sub>2</sub>' in place of 'CO'.  
(ii) The electrolyte was given wrongly by many candidates.
- (b) (i) The presence of unpaired d-electrons was not mentioned by many candidates.  
(ii) The polar nature of the x-y bonds or less overlapping of the orbitals was not mentioned by several candidates.  
(iii) The diamagnetic nature of  $\text{Cu}^+$  was not mentioned by several candidates.

### Suggestions for teachers

- Electrorefining should be taught with correct electrodes and electrolytes.
- The properties of transition metals should be explained with reference to the d-electrons.
- Magnetic nature should be explained in terms of electronic configuration.

### MARKING SCHEME

#### Question 7.

- (a) (i)  $\text{ZnO} + \text{C} \xrightarrow{1673\text{K}} \text{Zn} + \text{CO}$
- (ii) Anode – Impure Zn block  
Cathode – Thin sheet of pure Zinc  
Electrolyte –  $\text{ZnSO}_4$  solution containing a little dilute  $\text{H}_2\text{SO}_4$   
Cathode –  $\text{Zn}^{2+} + 2\text{e}^- \rightarrow \text{Zn}$   
Anode –  $\text{Zn} - 2\text{e}^- \rightarrow \text{Zn}^{2+}$
- (b) (i) Presence of unpaired electrons in d orbitals/ d-d transition  
Electrons absorb radiations of one colour from white light for excitation from one energy level to another within the same d subshell. Hence, transmitted light appears coloured.  
Or any correct explanation
- (ii) x – y bonds are weaker than x – x or y – y bonds.  
Polar nature of x-y bond.  
Less overlapping of orbitals in x – y bond.
- (iii)  $\text{Cu}^{+1}$  has no unpaired electrons, hence diamagnetic  
 $\text{Cu}^{+2}$  has unpaired electrons, hence paramagnetic

## SECTION C

*Answer any two questions.*

### Question 8


(a) How can the following conversions be brought about:

(i) Nitro benzene to benzene diazoniumchloride. [3]

(ii) Propanoic acid to ethylamine. [2]

(iii) Benzoic acid to benzaldehyde. [2]

(b) Identify the compounds A, B, C, D, E and F: [3]



#### Comments of Examiners



- (a) (i) Many candidates used  $\text{LiAlH}_4$  or  $\text{H}_2/\text{Ni}$  for the reduction of nitrobenzene to aniline. In some cases, reagents and temperature were not mentioned for diazotization.
- (ii) Several candidates started the conversion from ethanoic acid in place of propanoic acid. Conversion steps were incorrect in many cases.
- (iii) Most of the candidates reduced benzoic acid to benzaldehyde directly by metal / acid.
- (b) [E] was identified as methylamine and [F] as ethylacetate by many candidates.

#### Suggestions for teachers

- Reduction of nitrobenzene should be taught with correct reducing agent. Diazotization should be taught with correct reagents and temperature ( $0^\circ\text{C} - 5^\circ\text{C}$ ).
- Aliphatic and aromatic conversions should be practiced in class with correct reagents.
- Reactions should be practiced in class with correct reactants and products.

### MARKING SCHEME

#### Question 8.



- (b) [A] → Acetaldehyde or  $\text{CH}_3\text{CHO}$   
 [B] → Acetic acid or  $\text{CH}_3\text{COOH}$   
 [C] → Acetyl chloride or  $\text{CH}_3\text{COCl}$   
 [D] → Acetic anhydride or  $(\text{CH}_3\text{CO})_2\text{O}$   
 [E] → Acetaldoxime or  $\text{CH}_3\text{CH}=\text{NOH}$   
 [F] → Phenyl acetate or  $\text{C}_6\text{H}_5\text{OCOCH}_3$   
 OR any common name

### Question 9

- (a) Write balanced chemical equations for the following reactions and name the reactions: [3]  
 (i) Acetamide is heated with bromine and sodium hydroxide solution.  
 (ii) Benzaldehyde is treated with 50% sodium hydroxide solution.
- (b) Give one chemical test to distinguish between the following pairs of compounds: [3]  
 (i) Acetone and phenol.  
 (ii) Formic acid and Acetic acid.
- (c) (i) Name the type of isomerism exhibited by the following pairs of compounds: [2]  
 (1)  $(\text{C}_2\text{H}_5)_2\text{NH}$  and  $\text{CH}_3\text{-NH-C}_3\text{H}_7$   
 (2) 1-butanol and 2 methyl-1-propanol
- (ii) Name the type of isomerism that the compound with molecular formula  $\text{C}_3\text{H}_6\text{O}_2$  exhibits. Represent the isomers. [2]

### Comments of Examiners


- (a) (i) Only the main product was written by many candidates while the by products were missing. In some cases, the equation was not balanced.  
 (ii) The name of the reaction was given incorrectly by a number of candidates.
- (b) (i) Many candidates gave ferric chloride test for an acetone. Some only mentioned the name of the test without giving the observation.  
 (ii) Incorrect tests were given by many candidates.
- (c) (i) Some candidates gave the answer as 'position isomers'.  
 (ii) Most of the candidates were able to attempt this part correctly.

### *Suggestions for teachers*


- Name equations should be taught in balanced form. Reactants and products should be correctly taught.
- Identification of organic compounds should be done in class with correct tests.
- Isomerism should be explained by drawing the structural formulae of the isomers.

## MARKING SCHEME

### Question 9.



Hoffmann's bromanide or Hofmann's degradation reaction.




(b)(i) Acetone when treated with  $\text{I}_2 + \text{NaOH}$  gives yellow ppt. of  $\text{CHI}_3$  but phenol with  $\text{I}_2 + \text{NaOH}$  gives no such ppt.

Phenol with neutral  $\text{FeCl}_3$  solution gives violet colouration but Acetone gives no such observation (Or any correct test)


(ii) Formic acid gives white or grey ppt. with  $\text{HgCl}_2$  solution but Acetic acid does not give such observation (Or any correct test)

(c) (i) (1) Metamerism

(2) Chain isomerism



Optical Isomerism:



### Question 10

(a) Write balanced chemical equations for the following reactions: [4]

- Oxalic acid is treated with acidified potassium permanganate solution.
- Benzoic acid is treated with a mixture of concentrated nitric acid and concentrated sulphuric acid.
- Methyl magnesium iodide is treated with carbon dioxide and the product hydrolysed in acidic medium.
- Ethylacetate is treated with ammonia.



- (b) An organic compound [A] having molecular formula  $C_2H_7N$  on treatment with nitrous acid gives a compound [B] having molecular formula  $C_2H_6O$ . [B] on treatment with an organic compound [C] gives a carboxylic acid [D] and a sweet smelling compound [E]. Oxidation of [B] with acidified potassium dichromate also gives [D]. [4]
- (i) Identify [A], [B], [C], [D] and [E].
- (ii) Write balanced chemical equation of [D] with chlorine in the presence of red phosphorus and name the reaction.
- (c) Acetamide is amphoteric in nature. Give two equations to support this statement. [2]

#### Comments of Examiners


- (a) (i) A number of candidates wrote incorrect equations.
- (ii) Ortho - para benzoic acid was written as the product by several candidates.
- (iii) Some candidates wrote wrong products in this part.
- (iv) This part was generally done correctly by candidates.
- (b) (i) Identification of [C] was incorrectly done by several candidates.
- (ii) A number of candidates wrote unbalanced equations.
- (c) Many candidates did not attempt this part of the question. Some gave the reaction with NaOH to show the acidic nature of acetamide.

#### Suggestions for teachers

- Equation should be studied by writing the reactants and products with balanced form.
- Organic problems of identification should be given in class.
- The amphoteric nature of acetamide should be taught correctly with proper reactants and products.

#### MARKING SCHEME

##### Question 10.

- (a) (i) 
$$\begin{array}{c} 5\text{COOH} \\ | \\ \text{COOH} \end{array} + 2\text{KMnO}_4 + 3\text{H}_2\text{SO}_4 \rightarrow 10\text{CO}_2 + \text{K}_2\text{SO}_4 + 2\text{MnSO}_4 + 8\text{H}_2\text{O}$$
- (ii) 
- (iii) 
$$\text{CH}_3\text{MgI} + \text{O}=\text{C}=\text{O} \rightarrow [\text{IMgO} - \begin{array}{c} \text{CH}_3 \\ | \\ \text{C} \end{array} = \text{O}] \rightarrow \text{CH}_3\text{COOH} + \text{MgI}(\text{OH})$$
- (iv) 
$$\text{CH}_3\text{COOC}_2\text{H}_5 + \text{NH}_3 \rightarrow \text{CH}_3\text{CONH}_2 + \text{C}_2\text{H}_5\text{OH}$$

- (b) (i) [A]  $\rightarrow$   $\text{CH}_3\text{CH}_2\text{NH}_2$  or Ethylamine  
 [B]  $\rightarrow$   $\text{CH}_3\text{CH}_2\text{OH}$  or Ethanol  
 [C]  $\rightarrow$   $(\text{CH}_3\text{CO})_2\text{O}$  or Ethanoic anhydride  
 [D]  $\rightarrow$   $\text{CH}_3\text{COOH}$  or Ethanoic acid.  
 [E]  $\rightarrow$   $\text{CH}_3\text{COOC}_2\text{H}_5$  or Ethyl ethanoate.
- (ii)  $\text{CH}_3\text{COOH} + \text{Cl}_2 \xrightarrow{\text{Red P}} \text{CH}_2\text{ClCOOH} + \text{HCl}$
- Hell Volhard Zelinsky reaction or  
 HVZ reaction.
- (c) Basic :  $\text{CH}_3\text{CONH}_2 + \text{HCl} \rightarrow \text{CH}_3\text{CONH}_2 \cdot \text{HCl}$  or  $[\text{CH}_3\text{CONH}_3]^+\text{Cl}^-$   
 Acidic:  $\text{CH}_3\text{CONH}_2 + \text{HgO} \rightarrow (\text{CH}_3\text{CONH})_2\text{Hg} + \text{H}_2\text{O}$

### **GENERAL COMMENTS:**

#### **(a) Topics found difficult by candidates in the Question Paper:**

- Co-ordination compounds.
- Definitions of different terms and units.
- Isomerism.
- Buffer solution
- Electrochemical cells.
- Geometry and hybridization.
- Inorganic equations.

#### **(b) Concepts between which candidates got confused:**

- Hydrolysis of salts.
- Catalysts in organic reactions.
- Conversion in Organic Chemistry.
- Numericals in Colligative properties with correct units.

#### **(c) Suggestions for students:**

- Practice organic conversions.
- Study chemical tests to distinguish between organic compounds.
- Study isomerism by drawing the structural formulae of the isomers.
- Avoid selective studies.
- Practice equations, both inorganic and organic with balanced form and correct condition.
- Practice numericals from all chapters with correct formulae and units.